

University of Groningen

Dark-Matter in Galaxies from Gravitational Lensing and Stellar Dynamics Studies

Koopmans, L. V. E.

Published in:
 Highlights of Astronomy

DOI:
[10.1017/S1743921310008276](https://doi.org/10.1017/S1743921310008276)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Koopmans, L. V. E. (2010). Dark-Matter in Galaxies from Gravitational Lensing and Stellar Dynamics Studies. In IF. Corbett (Ed.), *Highlights of Astronomy* (Vol. 15, pp. 74). (IAU Symposium Proceedings Series; Vol. 5, No. H15). Cambridge University Press. <https://doi.org/10.1017/S1743921310008276>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Dark-Matter in Galaxies from Gravitational Lensing and Stellar Dynamics Studies

L. V. E. Koopmans

Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700AV, Groningen, The Netherlands; email: koopmans@astro.rug.nl

Abstract. Strong gravitational lensing and stellar dynamics provide two complementary methods in the study of the mass distribution of dark matter in galaxies out to redshift of unity. They are particularly powerful in the determination of the total mass and the density profile of mass early-type galaxies on kpc to tens of kpc scales, and also reveal the presence of mass-substructure on sub-kpc scale. I will shortly discuss these topics in this review.

Keywords. Galaxies: elliptical and lenticular, cD – Galaxies: formation – Galaxies: evolution

In recent years, the combined study of early-type galaxies using stellar dynamics and gravitational lensing has born exiting and unique results on the density profile of early type galaxies to redshift of unity, showing that massive early-type galaxies are close to homologous, have total density profile close to isothermal and do not appreciably evolve in their ensemble properties (for example Treu & Koopmans (2004), Koopmans *et al.* (2009)). Neither through lensing alone, nor through dynamics would these results have been possible to obtain. This has become even more exciting in recent years through the discovery of nearly 100 new strong galaxy-scale lens systems in the SLACS survey, for example Bolton *et al.* (2008) and development of powerful modeling techniques that self-consistently combine these two methods, using HST imaging and IFU/Longslit spectroscopic data Barnabè & Koopmans (2007), Czoske *et al.* (2008), Barnabè *et al.* (2009).

A second exiting new development is the *imaging* of mass-substructure on sub-kpc scales in these galaxies using extended arcs and Einstein rings Vegetti & Koopmans (2009). Also this is now possible thanks to the typical extended and highly-magnified nature of the lensed images in SLACS systems. Because multiple images allow one to separate the source structure from the lens-potential structure, this provides a unique method to probe unseen structure in lenses from dwarf satellites to general deviations from simple symmetries. Recently, the first detection of a high M/L-ratio dwarf satellite was reported using this method. Detection (even non-detections) promise to be a unique avenue to study the low-mass end of the dwarf-satellite mass function and possibly even probe the nature of dark matter – Vegetti & Koopmans (2009).

References

- Barnabè, M. & Koopmans, L. V. E. 2007, ApJ, 666, 726
Barnabè, *et al.*, 2009, MNRAS, 399, 21
Bolton, A. S., *et al.* 2008, ApJ, 682, 964
Czoske, O., *et al.* 2008, MNRAS, 384, 987
Koopmans, L. V. E., *et al.* 2009, ApJL, 703, L51
Treu, T. & Koopmans, L. V. E. 2004, ApJ, 611, 739
Vegetti, S. & Koopmans, L. V. E. 2009, MNRAS, 1456
Vegetti, S., Koopmans, L. V. E., Bolton, A., Treu, T., & Gavazzi, R. 2009, arXiv:0910.0760
Vegetti, S. & Koopmans, L. V. E. 2009, MNRAS, 392, 945